

THE MAX NOETHER FUNDAMENTAL THEOREM IS COMBINATORIAL

J.I. COGOLLUDO AGUSTÍN AND M.Á.MARCO BUZUNÁRIZ

ABSTRACT. In the present paper we give a reformulation of the Noether Fundamental Theorem for the special case where the three curves involved have the same degree. In this reformulation, the local Noether's Conditions are weakened. To do so we introduce the concept of Abstract Curve Combinatorics (ACC) which will be, in the context of plane curves, the analogue of matroids for hyperplane arrangements.

1. INTRODUCTION

In 1873 Noether stated his celebrated Fundamental Theorem [8], sometimes referred to as the “ $AF + BG$ ” Theorem. This theorem brings together the geometric and algebraic conditions plane projective algebraic curves should satisfy when belonging to a pencil. The following statement can be found in [3].

Theorem 1.1 (Max Noether's Fundamental Theorem). *Let F, G, H be homogeneous reduced polynomials in three variables defining projective algebraic curves $V(F)$, $V(G)$, and $V(H)$. Assume $V(F)$ and $V(G)$ have no common components. Then there is an equation $H = AF + BG$ (with A, B forms of degrees $\deg(H) - \deg(F)$ and $\deg(H) - \deg(G)$ respectively) if and only if $H_P \in (F_P, G_P) \subset \mathcal{O}_P(\mathbb{P}^2)$ for any $P \in V(F) \cap V(G)$.*

Here we denote by F_P the germ of F at P , by (F_P, G_P) the local ideal generated by the germs F_P and G_P , and by $V(F) \subset \mathbb{P}^2$ the set of zeroes of F . The local conditions on the equations F, G, H are called *Noether conditions*.

This theorem was originally attacked both from geometric and algebraic points of view ([9, 3]) and it has been recently generalized to the non-reduced case by Fulton [4].

Most of the efforts to understand and rewrite Noether's Fundamental Theorem have been focused on finding conditions that are equivalent to the Noether conditions in particular instances like transversality of branches, ordinary singularities, etc.

Our purpose here is to concentrate on the case where $\deg F = \deg G = \deg H$ and to weaken the Noether's conditions so as to have strictly weaker local conditions that can still provide the equivalence of the result. Note that the Noether Fundamental Theorem is a combination of a global condition (the existence of the curves F, G, H) and local conditions. Our weakened local conditions combined with the global condition result in this equivalence.

The weakened local conditions can be briefly described as follows: We say F satisfies the *combinatorial conditions* with respect to G and H if for any point $P \in V(F) \cap V(GH)$ and any local branch δ of F at P then $\mu_P(\delta, G) = \mu_P(\delta, H)$, where μ_P denotes the multiplicity of intersection of branches at P . Also we say that F, G, H satisfy the conditions for a *combinatorial pencil* if each equation satisfies the combinatorial conditions with respect to the other two

2000 *Mathematics Subject Classification.* 58K65, 14Q05, 14B05, 14C22, 14H50, 14F25, 32K07.

Key words and phrases. Plane algebraic curves, singularities.

equations. We also introduce the concept of a *primitive combinatorial pencil* which corresponds with the geometric idea that the fibers of the map over $\mathbb{P}^1$ induced by the pencil after resolution of indeterminacy are connected. In §3 we prove that any combinatorial pencil can be refined to a primitive combinatorial pencil.

The global condition can be rewritten as follows: If $\deg F = \deg G = \deg H$, then the condition $H = AF + BG$ simply means that H belongs to the pencil generated by F and G , or simply that F, G, H belong to a pencil.

The main result is the following.

Theorem 1.2. *Let F, G, H be projective plane curves of the same degree. Assume F and G have no common components. If F, G, H belong to a primitive combinatorial pencil then they belong to a pencil.*

To end this introduction we present two examples aimed to clarify the sharpness of these combinatorial conditions. The first one points out that the combinatorial conditions are indeed weaker than the Noether conditions and the second one suggests that the conditions cannot be weakened.

Example 1.3. This first example shows that the (local) Noether Conditions are stronger than the combinatorial condition described above. Consider the germs $f = x^3$, $g = y^2$ and $h = y^2 + (x+y)^3$ in $\mathcal{O}_P(\mathbb{P}^2) = \mathbb{C}\{x, y\}$, $P = [0 : 0 : 1]$. It is obvious that they satisfy the combinatorial conditions at P since $\mu_P(f, g) = \mu_P(f, h) = \mu_P(g, h) = 6$. However, $h \notin (f, g)$ since $h = y^2 + x^3 + y^3 + 3xy^2 + 3x^2y$, where $h_1 = y^2 + x^3 + y^3 + 3xy^2 \in (x^3, y^2)$, but $h_2 = x^2y \notin (x^3, y^2)$, and $h = h_1 + h_2$.

Example 1.4. This second example shows that the combinatorial conditions have to be stated for each branch, as opposed to each irreducible component. Consider $F = ZY^2 - ZX^2 + X^3$, $G = (X + Y)^3$, and $H = (X - Y)^3$, three cubics. Note that $V(F) \cap V(G) = V(F) \cap V(H) = V(G) \cap V(H) = \{P = [0 : 0 : 1]\}$ and $\mu_P(F, G) = \mu_P(F, H) = \mu_P(G, H) = 9$. However, F, G, H are not in a pencil. Note that the combinatorial conditions are *not* satisfied, since F is not locally irreducible at P and the two branches δ_1 and δ_2 satisfy $\mu_P(\delta_1, G) = \mu_P(\delta_2, H) = 6$, and $\mu_P(\delta_1, H) = \mu_P(\delta_2, G) = 3$.

2. SETTINGS

2.1. Abstract Curve Combinatorics.

Definition 2.1. An *Abstract Curve Combinatorics* (ACC for short) is a sextuplet $W := (\mathbf{r}, S, \Delta, \partial, \phi, \mu)$, where

- (1) $\mathbf{r}$, S , and Δ are finite sets,
- (2) $\partial : \Delta \rightarrow S$ and $\phi : \Delta \rightarrow \mathbf{r}$ are surjective maps,
- (3) $\mu : \text{SP}^2(\Delta) \rightarrow \mathbb{N}$, where $\text{SP}^2(\Delta)$ is the symmetric product of Δ , such that $\mu(\delta_1, \delta_2) > 0$ if and only if $\partial(\delta_1) = \partial(\delta_2)$ and $\phi(\delta_1) \neq \phi(\delta_2)$.

For simplicity, we denote $\Delta_P := \partial^{-1}(P)$, $P \in S$.

We say that two ACC's are equivalent if there are bijections preserving the corresponding maps.

Remark 2.2. Note that any projective curve $\mathcal{C} \subset \mathbb{P}^2$ determines naturally an ACC $W_{\mathcal{C}} := (\mathbf{r}, S, \Delta, \partial, \phi, \mu)$, (which will be referred to as the *Weak Combinatorial Type of $\mathcal{C}$*) as follows:

- (i) The set $\mathbf{r}$ is the set of irreducible components of $\mathcal{C}$,
- (ii) The set $S := \text{Sing}(\mathcal{C})$, is the set of singular points of $\mathcal{C}$,
- (iii) $\Delta := \cup_{P \in S} \{\Delta_P\}$ where Δ_P is the set of local branches of $\mathcal{C}$ at $P \in S$, $\partial(\delta) := P$ if $\delta \in \Delta_P$, and ϕ assigns to each local branch the global irreducible component that contains it.
- (iv) $\mu(\delta_1, \delta_2)$ is defined as the multiplicity of intersection between δ_1 and δ_2 (when $\partial(\delta_1) = \partial(\delta_2)$ and $\phi(\delta_1) \neq \phi(\delta_2)$) and as zero otherwise.

In accordance with this motivation, given an ACC $W = (\mathbf{r}, S, \Delta, \partial, \phi, \mu)$, we will refer to the elements of $\mathbf{r}$ (resp. S , and Δ) as *irreducible components*, (points, and branches). Also μ will be referred to as the *intersection multiplicity* of two branches.

2.2. Bézout Condition and degrees. Consider W an ACC and define $d_{i,j} := \sum_{\substack{\phi(\delta_1) = i, \\ \phi(\delta_2) = j}} \mu(\delta_1, \delta_2)$, for any $i, j \in \mathbf{r}, i \neq j$.

Definition 2.3. W satisfies the *Bézout Condition* if $\frac{d_{i,j}d_{i,k}}{d_{j,k}}$ is independent of $j, k \in \mathbf{r}$. In that case, one can define

$$d_i := +\sqrt{\frac{d_{i,j}d_{i,k}}{d_{j,k}}}.$$

and will be referred to as the *degree* of i .

Note that the Weak Combinatorial Type of a plane projective curve satisfies the Bézout Condition and d_i coincides with the algebraic degree of the irreducible component i .

2.3. Combinatorial Pencils. Let W be an ACC satisfying the Bézout Condition.

Definition 2.4. We say that W contains a combinatorial pencil if there exist $\bar{m} := (m_i)_{i \in \mathbf{r}}$ a list of integers and $\mathcal{F} = \{F_1, \dots, F_k\}$, $k \geq 3$ a partition of $\mathbf{r}$ such that:

- (1) $\sum_{i \in F_j} m_i d_i$ is independent of $j \in \{1, \dots, k\}$, (such constant will be denoted by $d_{\mathcal{F}}$) and
- (2) for any $P \in S$ one of the following two conditions is satisfied:
 - (a) either $\phi(\Delta_P) \subset F_i$ for a certain $i = 1, \dots, k$,
 - (b) or $\phi(\Delta_P) \not\subset F_i$, in which case for each $\delta \in \Delta_P$, the natural number

$$\sum_{\phi(\delta') \in F_j} m_{\phi(\delta')} \mu(\delta, \delta')$$

is independent of j (as long as $\phi(\delta) \notin F_j$). Such a constant will be denoted by k_{δ} .

The points $P \in S$ satisfying (2b) will be called the *base points of the combinatorial pencil* and each $F_i \in \mathcal{F}$ will be called a *fiber*. The integer m_i will be called the *multiplicity* of the i -th component and the members of the partition $\mathcal{F}$ are the *members* of the pencil.

We also say that three curves F , G and H belong to a combinatorial pencil if $(\{F, G, H\}, \bar{m})$ is a combinatorial pencil, where $\bar{m}$ is the list of multiplicities of the components of $\mathcal{C} = F \cup G \cup H$.

Our purpose will be to investigate under what circumstances three curves belonging to a combinatorial pencil, also belong to a pencil, that is $H = AF + BG$ for some $A, B \in \mathbb{C}^*$. Note that this is not true in general as one can simply see with line arrangements. Consider $F = XY$, $G = X^2 - Y^2$, and $H = X^2 - 4Y^2$. It is obvious that F , G , and H belong to a combinatorial pencil, but not to a pencil. The geometrical reason behind this phenomenon is that the resolution of the rational map $[X : Y : Z] \mapsto [F : G]$ does not have connected fibers.

One needs an extra condition that assures that the pencil is primitive. The definition of a primitive combinatorial pencil and the fact that any combinatorial pencil can be reduced to a primitive one will be the main idea of the coming section.

3. COMBINATORIAL VERSION OF THE STEIN FACTORIZATION THEOREM

In our context of pencils in $\mathbb{P}^2$, the Stein Factorization Theorem ([5, Corollary III.11.5]) and the fact that a rational surface is simply connected imply that any morphism f from a rational surface S onto $\mathbb{P}^1$ factorizes through a morphism $g : S \rightarrow \mathbb{P}^1$ with connected fibers and a covering c of $\mathbb{P}^1$, that is, $f = c \circ g$. In other words, any pencil whose resolution does not result into connected fibers can be refined (after a base change) into a pencil with connected fibers (also known as *primitive pencil*).

From a purely combinatorial point of view one can ask themselves if any combinatorial pencil admits a refinement into a primitive combinatorial pencil.

Similar results for line arrangements already exist (see [7, 2]).

3.1. Weak combinatorics of resolutions. In this section we will construct a combinatorial analogue of a blowing-up process, which will lead to the concept of solvable ACC. Such combinatorics have the appropriate structure for our purpose.

Definition 3.1. Let $W = (\mathbf{r}, S, \Delta, \partial, \phi, \mu)$ be an ACC. Let us fix a point $P \in S$ and a list $\bar{\nu} := (\nu_\delta)_{\delta \in \Delta_P}$ of positive integers. We say that the ACC $\widehat{W} = (\hat{\mathbf{r}}, \hat{S}, \hat{\Delta}, \hat{\partial}, \hat{\phi}, \hat{\mu})$ is obtained as a σ -process at P from W (denoted by $W \leftarrow \widehat{W}$) if there exists a partition $\{\hat{P}_1, \dots, \hat{P}_\ell\}$ of Δ_P such that the following properties are held:

- (1) $\hat{\mathbf{r}} = \mathbf{r} \cup \{E\}$ (intuitively, $\hat{\mathbf{r}}$ results from adding the *exceptional divisor* E to $\mathbf{r}$),
- (2) $\hat{S} = (S \setminus \{P\}) \cup \{\hat{P}_1, \dots, \hat{P}_\ell\}$, (the point P is replaced by the *infinitely near points* $\{\hat{P}_1, \dots, \hat{P}_\ell\}$),
- (3) $\hat{\Delta} = \Delta \cup \{\hat{\delta}_1, \dots, \hat{\delta}_\ell\}$, where $\hat{\phi}(\hat{\delta}_i) = E$, $\hat{\partial}(\hat{\delta}_i) = \hat{P}_i$, $\hat{\phi}|_\Delta = \phi$, $\hat{\partial}|_{(\Delta \setminus \Delta_P)} = \partial|_{(\Delta \setminus \Delta_P)}$, and $\hat{\partial}(\delta) = \hat{P}_i$ if $\delta \in \hat{P}_i$ (the exceptional divisor contributes with one local branch at each infinitely near point and the maps are naturally extended),
- (4) $\hat{\mu}(\delta, \hat{\delta}_i) = \nu_i$ if $\delta \in \hat{P}_i$ (this is the intersection multiplicity of each branch with the exceptional divisor),
- (5) $\hat{\mu}(\delta_1, \delta_2) = \mu(\delta_1, \delta_2) - \nu_{\delta_1} \nu_{\delta_2}$, if $\delta_1, \delta_2 \in \Delta_P$ and $\phi(\delta_1) \neq \phi(\delta_2)$, (the intersection multiplicity of two local branches after blow up decreases by the product of the multiplicities of the branches),
- (6) $\hat{\mu}$ extends μ outside $\text{SP}^2(\Delta_P)$.

Remark 3.2. Let W be the weak combinatorial type of a curve $\mathcal{C}$ in a rational surface V and let $V \leftarrow \hat{V}$ be a blow-up of V at a singular point P of $\mathcal{C}$. Note that then the weak combinatorial type of the total transform $\hat{\mathcal{C}}$ is obtained by a σ -process at P from W by using as $\bar{\nu}$ the list of multiplicities of the local branches at P .

This way one can extend the concept of resolution to general ACC's.

Definition 3.3. A sequence of σ -processes $W = W_0 \leftarrow W_1 \leftarrow \dots \leftarrow W_n$ of ACC's $W_k := (\mathbf{r}^{(k)}, S^{(k)}, \partial^{(k)}, \Delta^{(k)}, \phi^{(k)}, \mu^{(k)})$, $k = 0, 1, \dots, n$ is called a *resolution* of W if:

- (1) $\nu_\delta \leq 1$ if $\delta \notin \Delta^{(0)}$, where ν_δ is the multiplicity associated with δ at a σ -process, and

- (2) (Normal-crossing condition) $\#\Delta_P^{(n)} = 2$ for any $P \in S^{(n)}$ and $\mu^{(n)}$ only takes values in $\{0, 1\}$.

An ACC is called *solvable* if there exists a resolution.

Remark 3.4. Note that the ACC obtained from a curve in $\mathbb{P}^2$ admits a (combinatorial) resolution given by any (geometric) resolution of its singularities, that is, every weak combinatorial type is solvable. Such a resolution will be called a *geometric* resolution of W . Note that weak combinatorial types might admit non-geometric resolutions aswell.

3.2. Admissibility conditions. Let W be an ACC and let $\{v_i\}_{i \in \mathbf{r}}$ be a list of vectors in $\mathbb{K}^k$. For any $\delta \in \Delta$ define

$$(1) \quad v_\delta := \sum_{\substack{j \in \mathbf{r} \\ j \neq \phi(\delta)}} \mu(\delta, j) v_j,$$

where $\mu(\delta, j) := \sum_{\delta' \in \phi^{-1}(j)} \mu(\delta, \delta')$. Note that, by Definition 2.1.(3), the only branches that contribute to v_δ are those in Δ_P .

We say that $\{v_i\}_{i \in \mathbf{r}}$ satisfies the *admissibility conditions* for W if:

$$(2) \quad \{v_{\phi(\delta)}, v_\delta\}, \quad \text{are linearly dependent for all } \delta \in \Delta.$$

We will often denote this by saying $v_{\phi(\delta)} || v_\delta$ (note that one of the vectors might be zero).

Definition 3.5. A list of vectors $v_W := \{v_i\}_{i \in \mathbf{r}}$ in $\mathbb{K}^k$ satisfying the admissibility conditions (2) for W and spanning $\mathbb{K}^k$ is called a *k-admissible family* for W .

One has the following result:

Proposition 3.6. *If $(\mathcal{F}, \bar{m})$ is a combinatorial pencil of $(k+1)$ -fibers of W , then there exists a k-admissible family for W .*

Proof. Let us consider $\mathcal{F} = \{F_0, F_1, \dots, F_k\}$ and define the following family of vectors $v_i, i \in \mathbf{r}$:

$$v_i := \begin{cases} m_i e_j & \text{if } i \in F_j, j \neq 0 \\ -m_i(e_1 + \dots + e_k) & \text{if } i \in F_0. \end{cases}$$

Under these conditions note that if $P \in S$ is not a base point, then condition (2) is immediately satisfied since all the vectors involved are linearly dependent. Now, if $P \in S$ is a base point, then the condition (2b) in Definition 2.4 above implies that $k_\delta v_i + v_\delta = 0$ and hence condition (2) is also true. $\square$

Definition 3.7. The *k-admissible family* for W associated with the combinatorial pencil $(\mathcal{F}, \bar{m})$ as in Proposition 3.6 will be referred to as *the admissible family of W associated with $(\mathcal{F}, \bar{m})$* .

We need the following result from linear algebra.

Lemma 3.8. *Suppose $v_1, \dots, v_r \in \mathbb{K}^k$ are vectors such that*

$$\begin{aligned} v_1 &|| \sum_{j=1}^r a_{1,j} v_j, \\ v_2 &|| \sum_{j=1}^r a_{2,j} v_j, \\ &\dots \\ v_{r-1} &|| \sum_{j=1}^r a_{r-1,j} v_j, \end{aligned}$$

where $a_{i,j} = a_{j,i}$. Then

$$v_r \parallel \sum_{j=1}^r a_{r,j} v_j.$$

Proof. Note that if $v_r = 0$ or $v_1 = v_2 = \dots = v_{r-1} = 0$, then the result is immediate. Also, if any of the vectors $\{v_i\}_{i=1,\dots,r-1}$ are trivial, say $v_1 = 0$, it is enough to solve the same problem on the remaining vectors, since the coefficients $a_{1,i} = a_{i,1}$ either multiply the vector v_1 (and hence they have no contribution) or multiply v_i in the condition $v_1 \parallel \sum_{j=1}^r a_{1,j} v_j$, which is trivially satisfied since $v_1 = 0$.

Therefore, we will assume that all the vectors v_i ($i = 1, \dots, r$) are non-zero. Note that if $v \neq 0$, then $v \parallel w$, implies the existence of $\lambda \in \mathbb{K}$ such that $\lambda v + w = 0$. Hence, in our case, there exist $\lambda_i \in \mathbb{K}$ such that

$$(3) \quad \lambda_i v_i + \sum_{j=1}^r a_{i,j} v_j = 0$$

for $i = 1, \dots, r-1$. Consider an $r \times r$ symmetric matrix $A := (\alpha_{i,j})$ where the first $r-1$ rows are given by the coefficients of the equations (3) over the variables v_i , and the last row is given by the coefficients of $\sum_{j=1}^{r-1} a_{r,j} v_j$ over the same variables. Also consider $V := (v_{i,j})$ an $r \times k$ matrix whose i -th row is given by the coefficients of v_i . By (3) one has that

$$AV = \begin{pmatrix} 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & 0 \\ & & \dots & & \\ 0 & 0 & \dots & 0 & 0 \\ b_1 & b_2 & \dots & b_{k-1} & b_k \end{pmatrix},$$

and hence

$$V^t AV = \bar{b}^t v_r,$$

where $\bar{b} = (b_1, b_2, \dots, b_{k-1}, b_k)$ (row notation) is the vector of coordinates of $\sum_{j=1}^{r-1} a_{r,j} v_j$. Since $V^t AV = \bar{b}^t v_r$ is symmetric, one obtains that $v_r \parallel \sum_{j=1}^{r-1} a_{r,j} v_j$ and therefore $v_r \parallel \sum_{j=1}^r a_{r,j} v_j$. $\square$

In other words, the admissibility conditions for each point are redundant.

In what follows we look into how the admissibility conditions change under a σ -process. Consider $v_W := (v_i)_{i \in \mathbf{r}}$ a k -admissible family of vectors for W , and $\widehat{W}$ a σ -process of W at P associated with the multiplicity list $\bar{\nu} := (\nu_\delta)_{\delta \in \Delta_P}$.

One has the following.

Proposition 3.9. *Let W be an ACC and $\widehat{W}$ a σ -process of W . Then any k -admissible family v_W on W induces a k -admissible family $v_{\widehat{W}}$ on $\widehat{W}$ and vice versa.*

Proof. Let v_W be an admissible family on W . Then $v_{\widehat{W}} = (\hat{v}_i)_{i \in \hat{\mathbf{r}}}$ is defined as follows: $\hat{v}_i = v_i$ if $i \in \mathbf{r}$. The new vector associated with the exceptional divisor $\hat{v}_E$ is defined as:

$$(4) \quad \hat{v}_E := \sum_{\delta' \in \Delta_P} \nu_{\delta'} v_{\phi(\delta')}.$$

It remains to verify that the new admissible conditions are satisfied. In order to avoid ambiguity, all the new vectors in $\widehat{W}$ will be denoted as $\hat{v}$. First we fix an infinitely near point $\hat{P}$, a branch $\delta \in \hat{\Delta}_{\hat{P}}$, and assume $i := \phi(\delta)$. We have two cases:

(1) If $\delta \in \Delta_P$, then

$$\hat{v}_\delta := \sum_{\substack{j \in \mathbf{f} \\ j \neq i}} \hat{\mu}(\delta, j) v_j = \sum_{\substack{j \in \mathbf{r} \\ j \neq i}} \left(\mu(\delta, j) - \nu_\delta \left(\sum_{\substack{\delta' \in \Delta_P \\ \phi(\delta') = j}} \nu_{\delta'} \right) \right) v_j + \nu_\delta \hat{v}_E,$$

(see (1)). Therefore

$$(5) \quad \hat{v}_\delta = v_\delta + \nu_\delta \left(\sum_{\substack{\delta' \in \Delta_P \\ \phi(\delta') = i}} \nu_{\delta'} \right) v_i.$$

By hypothesis $v_\delta || v_i$, thus $\hat{v}_\delta || v_i$.

(2) If $\delta = \hat{\delta}$ is the branch of the exceptional divisor E at $\hat{P}$. Since there is only one such branch at $\hat{P}$, then Lemma 3.8 and the previous case gives the result.

The converse is immediate since according to (5) $v_\delta || v_i$ if and only if $\hat{v}_\delta || v_i$. $\square$

3.3. Dicritical divisors and connectedness of an ACC.

Definition 3.10. Let v_W be a k -admissible family of vectors. Consider $\widehat{W}$ a resolution of W and $v_{\widehat{W}}$ the k -admissible family obtained from v_W as in Proposition 3.9. An exceptional divisor E is called a *dicritical divisor* in $\widehat{W}$ if there exist at least two branches δ_1 and δ_2 in $\hat{\Delta}$ such that $D_1 := \phi(\delta_1) \neq \phi(\delta_2) =: D_2$ and $\hat{\mu}(\delta_1, E) = \hat{\mu}(\delta_2, E) = 1$ and $\{v_{D_1}, v_{D_2}\}$ generate a two-dimensional space. Also, an exceptional divisor E is called *trivial* if $v_E = 0$.

Remark 3.11. Under the conditions above, any dicritical divisor is trivial. This is a direct consequence of the fact that $\widehat{W}$ must satisfy the admissibility conditions. Since $\widehat{W}$ is a resolution, one has that $v_{D_1} || v_E$ and $v_{D_2} || v_E$, and $v_{D_1} \not|| v_{D_2}$ which is only true if $v_E = 0$.

However, note that the converse is not true in general. For example, consider three smooth conics in a pencil of conics that are pairwise bitangent and use the vectors $v_1 = (1, 0)$, $v_2 = (0, 1)$, and $v_3 = (-1, -1)$ for each conic. After the first blowing up of any base point one obtains an exceptional divisor E which will not become a dicritical divisor, but whose associated vector v_E is zero, since $v_E = v_1 + v_2 + v_3$.

Definition 3.12. We say two components of an ACC *intersect*, if there are branches of each intersecting with positive multiplicity of intersection. Analogously, we say two components A , B are *connected* if there is a sequence of components intersecting pairwise. Formally, A , B are connected if there is a sequence of components $A_0 = A, A_1, \dots, A_n = B$, a sequence of branches $\delta_i, \delta'_i \in \Delta$, ($i = 1, \dots, n$) such that $\phi(\delta_i) = A_{i-1}$, $\phi(\delta'_i) = A_i$, satisfying $\mu(\delta_i, \delta'_i) \neq 0$. Therefore the concept of *connected components* of an ACC can be defined.

One has the following interesting result.

Lemma 3.13. *After resolution and after removing the trivial divisors, different fibers of a combinatorial pencil belong to different connected components.*

Proof. Let $(\mathcal{F}, \bar{m})$ be a combinatorial pencil in $\mathcal{C}$ and let v_W be its associated k -admissible family. Consider $W \leftarrow \widehat{W}$ a resolution of singularities of $\mathcal{C}$. By Proposition 3.9 the associated family of vectors $v_{\widehat{W}}$ also satisfies the admissibility conditions shown in (2). Since $\widehat{W}$ corresponds to the combinatorics of a normal crossing divisor the condition at each normal crossing of two divisors, say E and E' , means that either $v_E = kv_{E'}$, $k \in \mathbb{K}^*$, or $v_E = 0$, or $v_{E'} = 0$. In other words, the

vectors associated with $\widehat{W}$ are either multiples of the original vectors of W or zero. Moreover, after removing the trivial divisors, components from different fibers are disconnected. $\square$

3.4. Intersection matrix. Let W be a solvable ACC satisfying the Bézout conditions (Definition 2.3). Consider $W = W_0 \leftarrow W_1 \leftarrow \cdots \leftarrow W_n$ a resolution of W , and $v_W := (v_i)_{i \in \mathbf{r}}$ a k -admissible family of vectors. Denote by P_ℓ the point blown-up at each step $W_\ell \leftarrow W_{\ell+1}$ with multiplicity list $\bar{\nu}^{(\ell)} := (\nu_\delta^{(\ell)})_{\delta \in \Delta_P}$, and define by $\mathcal{P}$ the collection of such points. As seen in Proposition 3.9, associated with v_W there are k -admissible families $v_{W_\ell} := (v_i)_{i \in \mathbf{r}^{(\ell)}}$ of W_ℓ . Define $\tilde{\mathbf{r}}^{(n)} := \{i \in \mathbf{r}^{(n)} \mid v_i \neq 0\}$. Consider the following *incidence matrix* associated with the resolution W_n of W and with the admissible family v_W :

$$J := (a_{ij})_{i \in \tilde{\mathbf{r}}^{(n)}, j \in \mathcal{P}},$$

where

$$a_{ij} := \begin{cases} \nu_i^{(\ell)} & \text{if } j = P_\ell, i \in \mathbf{r}^{(\ell)} \\ -1 & \text{if } j = P_\ell, i = E_{\ell+1} \\ 0 & \text{otherwise,} \end{cases}$$

(recall that $\nu_i := \sum_{\substack{\delta \in \Delta_P \\ \phi(\delta) = i}} \nu_\delta$ and note that $\#\mathcal{P} = n$). Also define the *degree matrix* $D := \bar{d}^t \bar{d}$, where $\bar{d} := (d_i)_{i \in \tilde{\mathbf{r}}^{(n)}}$, and

$$d_i := \begin{cases} d_i & \text{if } i \in \mathbf{r}^{(0)} \\ 0 & \text{otherwise.} \end{cases}$$

Finally, we combine both matrices in order to define $Q := D - JJ^t$. Also, for convenience, if W is already normal crossing (2), then we set $JJ^t = 0$.

Note that Q is a square matrix of order $\#\tilde{\mathbf{r}}^{(n)}$.

Proposition 3.14. *If W satisfies the Bézout Conditions and $W \leftarrow W_n = \widehat{W} = (\hat{\mathbf{r}}, \hat{S}, \hat{\Delta}, \hat{\partial}, \hat{\phi}, \hat{\mu})$ is a resolution of W , then the matrix $Q = (q_{ij})_{i,j \in \tilde{\mathbf{r}}}$ satisfies the following:*

- (1) $q_{ij} = \hat{\mu}(i, j) := \sum_{\substack{\delta \in \hat{\phi}^{-1}(i) \\ \delta' \in \hat{\phi}^{-1}(j)}} \hat{\mu}(\delta, \delta')$ if $i \neq j$ (intuitively, the number of points at which the components i and j intersect in $\widehat{W}$). In particular $q_{ij} \geq 0$,
- (2) $q_{ii} = d_i^2 - \sum_{\ell \in L_i} (\nu_i^{(\ell)})^2$, where $L_i := \{\ell \mid i \in \phi^{(\ell)}(\Delta_{P_\ell}^{(\ell)})\}$.

Moreover, if the resolution is geometric, then Q is the intersection matrix of the non-trivial divisors of $\hat{\mathbf{r}}$ in $\widehat{W}$.

Proof. By definition of Q one has that

$$q_{ij} = d_i d_j - \sum_{\ell \in L_{ij}} \nu_i^{(\ell)} \nu_j^{(\ell)},$$

where $L_{ij} := \{\ell \mid i, j \in \phi^{(\ell)}(\Delta_{P_\ell}^{(\ell)})\}$. This implies (2).

By definition, one has $\mu^{(\ell)}(i, j) = \nu_i^{(\ell+1)} \nu_j^{(\ell+1)} + \mu^{(\ell+1)}(i, j)$. Also, if W satisfies the Bézout Conditions then $d_i d_j = \mu(i, j)$. Hence, $d_i d_j = \sum_{\ell \in L_{ij}} \nu_i^{(\ell)} \nu_j^{(\ell)} + \hat{\mu}(i, j)$ and (1) follows. Finally, since $\widehat{W}$ is a resolution, $q_{i,j}$ is exactly the number of points at which the components i and j intersect in $\widehat{W}$.

The second part is a consequence of the Noether formula for the multiplicity of intersection of branches after resolution. $\square$

Example 3.15. Consider the combinatorics corresponding to the arrangement of four curves: a smooth conic, two tangent lines to the conic, and the line joining the tangency points. We will order them and the exceptional divisors of the expected resolution as in Figure 1. The only non-




FIGURE 1. Resolution of a conic-line arrangement

trivial case corresponds to when E_3 and E_4 are the only dicritical divisors. The corresponding matrices follow:

$$J := \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 \end{bmatrix}, D := \begin{bmatrix} 4 & 2 & 2 & 2 & 0 & 0 \\ 2 & 1 & 1 & 1 & 0 & 0 \\ 2 & 1 & 1 & 1 & 0 & 0 \\ 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}, \text{ and } Q := \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 1 \\ 0 & 0 & 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 1 & 0 & -2 \end{bmatrix}.$$

Definition 3.16. Let $\mathcal{P}_1 := (\mathcal{F}_1, \bar{m}_1)$ and $\mathcal{P}_2 := (\mathcal{F}_2, \bar{m}_2)$ be combinatorial pencils of W . We say $\mathcal{P}_2$ is a *refinement* of $\mathcal{P}_1$ if the fibers of $\mathcal{P}_2$ are contained in the fibers of $\mathcal{P}_1$.

Definition 3.17. Let W be a solvable ACC and let $W \xleftarrow{\pi} \widehat{W}$ be a resolution, we say a combinatorial pencil $\mathcal{P} = (\mathcal{F}, \bar{m})$ is *primitive w.r.t. π* if each two components in the same fiber of $\mathcal{F}$ can be connected in $\widehat{W}$ by non-trivial divisors and the multiplicities of the components are coprime, that is, $\gcd(\bar{m}) = 1$.

If W is a weak curve combinatorics, then we simply call a combinatorial pencil *primitive* if it is primitive w.r.t. a geometric resolution.

Note that the combinatorial notion of primitive is again combinatorial, since the resolution is given again combinatorially.

Theorem 3.18. Let W be a solvable ACC satisfying Bézout Conditions and let $W \xleftarrow{\pi} \widehat{W}$ be a resolution, then any combinatorial pencil in W admits a primitive refinement w.r.t π .

Proof. According to Proposition 3.6 any combinatorial pencil admits a k -admissible family $v_W := \{v_i\}_{i \in \mathbf{r}}$. Take a resolution $W = W_0 \leftarrow W_1 \leftarrow \dots \leftarrow W_n = \widehat{W}$ and consider the resulting admissible family $v_{\widehat{W}}$.

By Lemma 3.13, there exist trivial divisors. After removing them, one can construct the matrices J , D and Q as above. Let A be the matrix whose columns are the non-zero vectors of $v_{\widehat{W}}$. It is easy to check that Bézout's Theorem implies $D \cdot A^t = 0$, and, by construction, $J^t \cdot A^t = 0$. So the rows of A are in the kernel of both Q and D . Note that the kernel of JJ^t coincides with the kernel of J^t .

After a suitable ordering of the elements of $\tilde{\mathbf{r}}^{(n)}$, one can assume that the matrix Q decomposes into a direct sum of irreducible boxes Q_λ , $\lambda = 1, \dots, \kappa$. This induces a partition $\tilde{\mathcal{F}} := \{\tilde{F}_1, \dots, \tilde{F}_\kappa\}$, (with $\tilde{F}_\lambda \subset \tilde{\mathbf{r}}^{(n)}$) of the components of the combinatorics, and hence, in the columns of A . Each submatrix Q_λ is symmetric, and by Proposition 3.14(1) it has non-negative entries outside the diagonal. Hence, using the Vinberg classification of matrices (see [6, Thm. 4.3]) on $-Q_\lambda$, one can ensure that Q_λ is of one of the following types:

- (Fin) $\det(Q_\lambda) \neq 0$; there exists a vector v with positive entries such that $Q_\lambda v$ has negative entries.
- (Aff) $\text{corank}(Q_\lambda) = 1$, and its kernel is generated by a vector with only positive entries.
- (Ind) There exists a vector u with only positive entries such that $Q_\lambda u$ has only positive entries.

Here (Fin), (Aff), and (Ind) stand for Finite, Affine, and Indefinite types respectively. We have seen that Q has a nontrivial kernel, so the Q_λ cannot be an (Fin)-matrix. If one of the Q_λ , say Q_1 , is an (Ind)-matrix, one can consider a vector u_1 with positive entries such that $Q_1 u_1$ has only positive entries. For the rest of the Q_λ , one can find vectors u_λ with only negative entries such that $Q_\lambda u_\lambda$ has only zero entries (if Q_λ is an (Aff)-matrix) or only negative entries (if Q_λ is an (Ind)-matrix). By multiplying the u_λ by suitable positive numbers, one can reconstruct a vector u such that $Du = 0$.

Now denoting by $(\cdot, \cdot)$ the standard scalar product:

$$0 \geq -(J^t u, J^t u) = (Qu, u) - (Du, u) = (Qu, u) = (Q_1 u_1, u_1) + \sum_{i \geq 2} (Q_\lambda u_\lambda, u_\lambda) \geq (Q_1 u_1, u_1) > 0,$$

which leads to contradiction. So we can conclude that all the Q_λ are (Aff)-matrices.

Note that, by Proposition 3.14, the partition $\tilde{\mathcal{F}}$ induced by the boxes Q_λ is equal to the partition given by the connected components in $\tilde{\mathbf{r}}^{(n)}$.

Note that a vector is in the kernel of Q if and only if it is made up of vectors that are in the kernel of the Q_λ 's. In particular, the kernel of Q has dimension equal to the number of irreducible boxes.

From now on, K_Q will denote the kernel of Q , and K_D will denote the kernel of D , the degree matrix. Let u_λ be a positive vector that generates the kernel of the box Q_λ , and $\tilde{u}_\lambda$ the vector of $\mathbb{K}^{r_n}$, $r_n := \#\tilde{\mathbf{r}}^{(n)}$ obtained from u_λ by completing the entries corresponding to the other boxes with zeroes. As mentioned above, $\{\tilde{u}_\lambda\}_{\lambda=1, \dots, \kappa}$ is a basis of K_Q . Since $D = \bar{d}^t \cdot \bar{d}$, thus $K_D = \ker \bar{d}$, and thus $\text{codim } K_D = 1$. Also, since K_Q has a set of non-negative vectors as a basis, one has that $K_Q \not\subset K_D$, and thus $\dim K_Q \cap K_D = \dim K_Q - 1$. By Bézout's Theorem $d := \bar{d} \cdot \tilde{u}_\lambda$ is independent of λ and hence $\{\tilde{u}_\lambda - \tilde{u}_\kappa\}_{\lambda=1, \dots, \kappa-1}$ is a basis of $K_Q \cap K_D$.

Consider N , the matrix whose rows are the family of vectors $\{\tilde{u}_\lambda - \tilde{u}_\kappa\}_{\lambda=1, \dots, \kappa-1}$. As we have seen before, the rows of A must be a linear combination of the rows of N .

Let us construct the family of vectors $w := \{w_i\}_{i \in \mathbf{r}^{(n)}}$ as follows:

- $w_i = 0$ if $i \in \mathbf{r}^{(n)}$ is a trivial divisor.
- Otherwise, w_i is equal to the corresponding column of N .

Lemma 3.19. *The set of vectors w is a $(\kappa - 2)$ -admissible family for $\widehat{W}$.*

Proof. Since $\widehat{W}$ is a normal crossing ACC, the admissibility conditions are verified if whenever two components, say i and j intersect, then $w_i || w_j$.

If one of them, say i , is a trivial divisor, then there is nothing to prove, since $w_i = 0$. If neither of them are trivial divisors, then $i, j \in F_\lambda$ for some λ in the partition $\tilde{\mathcal{F}}$. Therefore v_i (and also v_j) must be proportional to e_λ (the obvious vector of the canonical basis) when $\lambda \neq \kappa$ and to $(-1, \dots, -1)$ if $\lambda = \kappa$. Thus the result follows. $\square$

All this shows that $\text{rank } N = \dim(K_Q \cap K_D)$, and the columns of N are multiples of the following κ vectors: the $\kappa - 1$ vectors e_λ of the canonical basis of $\mathbb{Z}^{\kappa-1}$ and the vector $e_\kappa := (-1, \dots, -1)$. There is essentially only one linear relation among them, which is $\sum_{\lambda=1}^\kappa e_\lambda = 0$. Moreover, if $i \in F_\lambda$, then $v_i = m_i \cdot e_\lambda$, where m_i is a positive rational number. Note that, without loss of generality, after multiplication by a natural number (for instance the least common multiple of the denominators) one can assume that the m_i 's are integer numbers.

Let us show that $\mathcal{F} := \tilde{\mathcal{F}} \cap \mathbf{r}$ and $\bar{m} := (m_i)_{i \in \mathbf{r}}$ defines a combinatorial pencil of κ fibers. In order to do so we need to check both properties in Definition 2.4. Let $P \in S$ not satisfying Property 2.4(2a) and let $\delta \in \Delta_P$ a branch of i at P . Since v is an admissible family, $\sum_{j \neq i} \mu(\delta, j) v_j = \sum_{\lambda=1}^\kappa \left(\sum_{j \in F_\lambda} \mu(\delta, j) m_j \right) e_\lambda$ is proportional to v_i , which forces the sum $k_\delta := \sum_{j \in F_\lambda} \mu(\delta, j) m_j$ to be independent of λ . $\square$

Remark 3.20. By Proposition 3.14, in the case where the combinatorial pencil is realizable and the resolution is geometric one can prove Theorem 3.18 without using the Vinberg classification of matrices. According to Proposition 3.14 in this case Q_λ is the intersection matrix of the divisors in F_λ . Using Zariski's Lemma for fibrations of surfaces (see [1, Lemma III.8.2]) a divisor $D = \sum m_i C_i$, where the divisors C_i belong to the same fiber F_λ , satisfies $D^2 = \bar{m} Q_\lambda \bar{m}^t = 0$ if and only if $pD = qF_\lambda$ for $p, q \in \mathbb{Z} \setminus \{0\}$. In particular, for line arrangements this provides another proof of the analogous result given in [7, 2].

Example 3.21. Let us further analyze Example 3.15. The matrix Q has three boxes $Q_1 = (0)$,

$$Q_2 = \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix}, \text{ and } Q_3 = \begin{bmatrix} -1 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}. \text{ Therefore } v_1 = (1), v_2 = (1, 1), \text{ and } v_3 =$$

$(2, 1, 1)$ generate the kernels of Q_1 , Q_2 and Q_3 and $v = (1, 1, 1, 2, 1, 1) \in K_Q \cap K_D$ according to the notation above. In fact, if $\mathcal{F} := \{\{1\}, \{2, 3\}, \{4\}\}$ and $\bar{m} := (1, 1, 1, 2)$, then $(\mathcal{F}, \bar{m})$ is a combinatorial pencil, which corresponds to the geometric pencil generated by the conic and the two lines, which contains the third line twice.

4. COMBINATORIAL MAX NOETHER FUNDAMENTAL THEOREM

We are ready to prove the following version of the Max Noether Fundamental Theorem.

Theorem 4.1. *Let $V(F) = \{F = f_1^{\ell_1} \cdot \dots \cdot f_p^{\ell_p} = 0\}$, $V(G) = \{G = g_1^{m_1} \cdot \dots \cdot g_q^{m_q} = 0\}$ and $V(H) = \{H = h_1^{n_1} \cdot \dots \cdot h_r^{n_r} = 0\}$ be three curves. If F, G and H are in a primitive combinatorial pencil, then F, G and H are in a pencil.*

Proof. Denote by $\mathcal{C} \subset \mathbb{P}^2$ the union of $V(F)$, $V(G)$ and $V(H)$ and by S the set of base points of the combinatorial pencil. Let us assume that all irreducible components of H can be connected outside the combinatorial pencil. Consider a local branch γ of H at a point $P \in S$. Denote $k := \mu_P(\gamma, F) = \mu_P(\gamma, G)$. Note that generically, one has that $\mu_P(\gamma, \alpha F + \beta G) = k$, but there exists a point $[\alpha_\gamma : \beta_\gamma] \in \mathbb{P}^1$ such that $\mu_P(\gamma, \alpha_\gamma F + \beta_\gamma G) > k$. Let us denote by h_1 the irreducible

component containing γ . Note that,

$$\begin{cases} \mu_Q(h_1, \alpha_\gamma F + \beta_\gamma G) > \mu_Q(h_1, F) = \mu_Q(h_1, G) & \text{if } Q = P \\ \mu_Q(h_1, \alpha_\gamma F + \beta_\gamma G) \geq \mu_Q(h_1, F) = \mu_Q(h_1, G) & \text{otherwise.} \end{cases}$$

Therefore, by Bézout, $d \cdot \deg h_1 = \sum_{Q \in S} \mu_Q(h_1, \alpha_\gamma F + \beta_\gamma G) > \sum_{Q \in S} \mu_Q(h_1, F) = d \cdot \deg h_1$, which implies that $\alpha_\gamma F + \beta_\gamma G$ is a multiple of h_1 . Consider now a resolution of the pencil, obtained by blowing up the base points $\pi : X \rightarrow \mathbb{P}^2$. The morphism $f : X \rightarrow \mathbb{P}^1$ is now well defined on the rational surface X and if $P \notin S$ then $f(\pi^{-1}(P)) := [F(P) : G(P)]$. By hypothesis, the preimage of $V(H)$ outside the dicritical divisors is connected and hence f is constant on the strict transform of $V(H)$, for any $Q \in \pi^{-1}(V(H))$ one has that $f(Q) = [-\beta : \alpha] \in \mathbb{P}^1$. On the other hand we know that for any h_1 and h_2 components of H one has that $\alpha_1 F + \beta_1 G = h_1 u_1$ and $\alpha_2 F + \beta_2 G = h_2 u_2$.

Consider now P_1 (resp. P_2) a regular point of $V(h_1) \setminus S$ (resp. $V(h_2) \setminus S$) and $Q_i := \pi^{-1}(P_i)$. By the remarks in the previous paragraph, $f(Q_i) := [F(P_i) : G(P_i)] = [-\beta : \alpha]$. Therefore $[-\beta_1 : \alpha_1] = [-\beta_2 : \alpha_2] = [-\beta : \alpha]$, and hence $\alpha F + \beta G = H'K$, where $H' = h_1^{n'_1} \cdot h_2^{n'_2} \cdot \dots \cdot h_r^{n'_r}$. We will denote this by $H'K_H \in (F, G)$.

Let us denote by Q_1 (resp. Q_2) the intersection matrix associated with a resolution of $F \cdot G \cdot H$ (resp. $F \cdot G \cdot H' \cdot K$) that dominates both. According to Proposition 3.14, Q_1 and Q_2 have the following form

$$Q_1 = \begin{bmatrix} Q_F & 0 & 0 \\ 0 & Q_G & 0 \\ 0 & 0 & Q_H \end{bmatrix} \quad Q_2 = \begin{bmatrix} Q_F & 0 & 0 & 0 \\ 0 & Q_G & 0 & 0 \\ 0 & 0 & Q_{H'} & M \\ 0 & 0 & M & Q_K \end{bmatrix}$$

where $Q_H = Q_{H'}$. Note that the box Q_H has a kernel of dimension 1 and hence $M = 0$, or else, the box $\tilde{Q} := \begin{bmatrix} Q_H & M \\ M & Q_K \end{bmatrix}$ would have a vector of type $(v_H, 0) \in \ker \tilde{Q}$, which contradicts $\tilde{Q}$ being of Affine type. Therefore we can assume that $M = 0$, $H' = (H'')^q$, $H = (H'')^p$, and

$$Q_2 = \begin{bmatrix} Q_F & 0 & 0 & 0 \\ 0 & Q_G & 0 & 0 \\ 0 & 0 & Q_H & 0 \\ 0 & 0 & 0 & Q_K \end{bmatrix}$$

which implies that the preimage of H'' and K by the resolution π are disconnected outside the dicritical divisors. By the algebraic Stein Factorization Theorem one can find a refinement of the pencil $(F, G, H'K)$ into a pencil $(\tilde{F}, \tilde{G}, \tilde{H})$. Since the original pencil is primitive, one has that $\tilde{F}^{n_F} = F$, $\tilde{G}^{n_G} = G$, and $\tilde{H}^{n_{H'}} = H' = (H'')^q$, that is, there exists $n_{H''} = \frac{n_{H'}}{q} \in \mathbb{N}$ such that $\tilde{H}^{n_{H''}} = H''$. By the hypothesis on degrees, this implies that $n_F = n_G = pn_{H''}$. Since $\gcd(n_F, n_G, pn_{H''}) = 1$ by hypothesis, then $n_F = n_G = pn_{H''} = 1$. Thus, $H = H'$ and therefore $\alpha F + \beta G = H$.

□

As an immediate corollary one has the following.

Corollary 4.2. *Let F, G, H be three homogeneous polynomials of the same degree in three variables such that their zero sets $V(F), V(G)$, and $V(H)$ are three irreducible curves with no common components. If $V(F) \cap V(G) = V(F) \cap V(H) = V(G) \cap V(H) = \{P\}$ and F, G , and H are locally irreducible at P , then $H = \alpha F + \beta G$.*

REFERENCES

1. W. Barth, C. Peters, and A. Van de Ven, *Compact complex surfaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 4, Springer-Verlag, Berlin, 1984. MR MR749574 (86c:32026)
2. M. Falk and S. Yuzvinsky, *Multinets, resonance varieties, and pencils of plane curves*, Compos. Math. **143** (2007), no. 4, 1069–1088. MR MR2339840
3. W. Fulton, *Algebraic curves. An introduction to algebraic geometry*, W. A. Benjamin, Inc., New York–Amsterdam, 1969, Notes written with the collaboration of Richard Weiss, Mathematics Lecture Notes Series.
4. ———, *Adjoints and Max Noether’s Fundamentalsatz*, Algebra, arithmetic and geometry with applications (West Lafayette, IN, 2000), Springer, Berlin, 2004, pp. 301–313. MR MR2037097 (2004m:14060)
5. Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York, 1977, Graduate Texts in Mathematics, No. 52. MR MR0463157 (57 #3116)
6. V.G. Kac, *Infinite-dimensional Lie algebras*, third ed., Cambridge University Press, Cambridge, 1990.
7. M.Á. Marco Buzunáriz, *Resonance varieties, admissible line combinatorics, and combinatorial pencils*, Preprint available at [arXiv:math.CO/0505435](https://arxiv.org/abs/math/0505435), 2005.
8. M. Nöther, *Ueber einen Satz aus der Theorie der algebraischen Functionen*, Math. Ann. **6** (1873), no. 3, 351–359. MR MR1509826
9. C.A. Scott, *A proof of Noether’s fundamental theorem*, Mathematische Annalen **52** (1899), no. 4, 593–597.

DEPARTAMENTO DE MATEMÁTICAS, IUMA, UNIVERSIDAD DE ZARAGOZA, C/ PEDRO CERBUNA, 12, E-50009 ZARAGOZA, SPAIN

E-mail address: jicogo@unizar.es

ICMAT: CSIC-COMPLUTENSE-AUTONOMA-CARLOS III, DEPARTAMENTO DE ÁLGEBRA - FACULTAD DE CC. MATEMÁTICAS - PLAZA DE LAS CIENCIAS, 3, 28040 MADRID, SPAIN

E-mail address: mmarco@unizar.es